

PURE and APPLIED ANALYSIS

PAA

MAARTEN V. DE HOOP, JOONAS ILMAVIRTA,
MATTI LASSAS AND TEEMU SAKSALA

**A FOLIATED AND REVERSIBLE FINSLER MANIFOLD IS
DETERMINED BY ITS BROKEN SCATTERING RELATION**

A FOLIATED AND REVERSIBLE FINSLER MANIFOLD IS DETERMINED BY ITS BROKEN SCATTERING RELATION

MAARTEN V. DE HOOP, JOONAS ILMAVIRTA, MATTI LASSAS AND TEEMU SAKSALA

The broken scattering relation consists of the total lengths of broken geodesics that start from the boundary, change direction once inside the manifold, and propagate to the boundary. We show that if two reversible Finsler manifolds satisfying a convex foliation condition have the same broken scattering relation, then they are isometric. This implies that some anisotropic material parameters of the earth can be in principle reconstructed from single scattering measurements at the surface.

1. Introduction

The broken scattering relation of a Finsler manifold with boundary describes all the scenarios where a geodesic starts inward at the boundary, changes direction at some point in the interior, and makes its way back to the boundary. The objective in this paper is to reconstruct the Finsler manifold from such data, provided that some assumptions are met. We have two key assumptions. One is that the Finsler geometry is reversible, meaning that the Minkowski norm on each tangent space is symmetric ($|-v| = |v|$) or, equivalently, that the reverse of a geodesic is a geodesic. We also assume that the manifold has a strictly convex foliation with a family of smooth hypersurfaces. In this paper we will also describe an important relationship between Finsler geometry and elastic waves.

Statement of the result. Let M be a smooth, compact manifold of dimension 3 or higher, with smooth boundary ∂M . We use the notation TM for the tangent bundle of M . Recall that a Finsler metric $F : TM \rightarrow \mathbb{R}$ is a continuous positive function such that on each fiber $T_x M$ of TM the function F is a Minkowski norm meaning:

- F is smooth outside the zero section.
- F is positively homogeneous of order 1; that is, $F(x, av) = aF(x, v)$ for any $a > 0$, $x \in M$ and $v \in T_x M$.
- F is convex in the sense that the local Riemannian metric

$$g_{ij}(x, v) := \frac{1}{2} \frac{\partial^2}{\partial v_i \partial v_j} [F^2](x, v), \quad i, j \in \{1, \dots, \dim M\},$$

is positive definite for any $(x, v) \in TM$, $v \neq 0$.

If F is a Finsler metric, we call the pair (M, F) a Finsler manifold. If the Finsler metric is symmetric with respect to the directional variable v in the sense of $F(x, v) = F(x, -v)$, we call F *reversible* Finsler metric.

MSC2020: 86A22, 53Z05, 53C60.

Keywords: Finsler manifold, scattering relation, distance functions, anisotropic elasticity.

Definition 1. A Finsler manifold with boundary is said to have a strictly convex foliation if there is a smooth function $f : M \rightarrow \mathbb{R}$ so that

- (i) $f^{-1}(0) = \partial M$, $f^{-1}(0, S) = \text{int}(M)$, $f^{-1}(S)$ has empty interior,
- (ii) for each $s \in [0, S)$ the set $\Sigma_s := f^{-1}(s)$ is a strictly convex smooth surface in the sense that $df \neq 0$ and any geodesic γ , having initial conditions in $T\Sigma_s$, satisfies $\partial_t^2 f(\gamma(t))|_{t=0} < 0$.

In a similar fashion, we may define a strictly convex surface without a foliation. For any hypersurface Σ there is a real-valued function f defined in its neighborhood so that $f = 0$ and $df \neq 0$ on Σ . The surface is called strictly convex if any geodesic $\gamma(t)$ tangent to Σ at $t = 0$ satisfies $\partial_t^2 f(\gamma(t))|_{t=0} < 0$.

In particular we say that a reversible Finsler manifold (M, F) has a strictly convex boundary ∂M if

$$f(x) = d(x, \partial M), \quad x \in U,$$

satisfies the former conditions near the boundary. In this case any pair of points in M can be connected by a distance-minimizing geodesic [Bartolo et al. 2011].

A manifold with a strictly convex foliation enjoys a certain weak nontrapping condition given in Lemma 14. If there is a trapped geodesic, it has to be asymptotically in the set $f^{-1}(S)$ where the foliating function f attains its maximum value, and it depends on the geometry of this set whether this is at all possible.

Examples of manifolds satisfying Definition 1 but having trapped geodesics can be constructed as follows. The following examples are Riemannian, but they can be modified to Finsler manifolds by varying the Riemannian Finsler function slightly in directions that are irrelevant for the definition. Take any smooth and symmetric function $\phi : [-1, 1] \rightarrow (0, \infty)$ and consider the surface of revolution

$$M_\phi = \{(x, y) \in \mathbb{R}^2 \times [-1, 1] : |x| = \phi(y)\} \subset \mathbb{R}^3$$

with the inherited Riemannian metric. If $\phi'' > 0$ on the whole interval, then the restriction $f|_{M_\phi} : M_\phi \rightarrow \mathbb{R}$ of $f(x, y) = 1 - y^2$ is a valid foliation function — this is evident upon inspection of the geodesic equation on a surface of revolution. See for instance [do Carmo 1992, Chapter 3, Exercise 1]. Now $S = 1$, and the set $f^{-1}(1)$ is the circle around the narrowest point of the surface and is therefore a trapped geodesic. There are also geodesics that are asymptotic to this trapped geodesic and are therefore also trapped; they can be produced by aiming a sequence of geodesics from a single point so that their closest point to the set $f^{-1}(1)$ gets smaller, and taking the limiting direction. A noncompact example is given by a half

$$M_\phi^+ = \{(x, y) \in M_\phi : y > 0\}$$

of the set M_ϕ defined above. It satisfies our convex foliation condition as well and has these trapped geodesics but $f^{-1}(1) = \emptyset$. (Compare this to the weak nontrapping property of Lemma 14.) If ϕ is the hyperbolic cosine, the manifold M_ϕ is the catenoid.

Next we describe what it means for two manifolds to have the same boundary data. The Finsler function $F : TM \rightarrow \mathbb{R}$ describes the geometry, and boundary information requires knowledge of $F|_{\partial TM}$. If we know the geometry of ∂M , we can deduce $F|_{T\partial M}$. However, $T\partial M \subset \partial TM$ is a subbundle with fiberwise codimension 1. In Riemannian geometry the knowledge of the metric on $T\partial M$ determines

uniquely the metric on ∂TM in boundary normal coordinates. In Finsler geometry it does not, because the geometry does not factorize as a product of tangential and normal directions of the boundary. Also a diffeomorphism $\xi : \partial M_1 \rightarrow \partial M_2$ only induces a diffeomorphism $d\xi : T \partial M_1 \rightarrow T \partial M_2$, and that is not enough for our purpose. Thus we introduce the following compatibility condition.

Definition 2. Let M_1 and M_2 be two smooth manifolds with boundary. We say that a diffeomorphism $\Xi : \partial TM_1 \rightarrow \partial TM_2$ is compatible with a diffeomorphism $\xi : \partial M_1 \rightarrow \partial M_2$ if Ξ is a linear isomorphism on every fiber and satisfies $\Xi(T \partial M_1) = T \partial M_2$ and $\Xi|_{T \partial M_1} = d\xi$.

Before showing that the broken scattering relation determines a Finsler manifold, we define what the relation is. To this end, let $\phi_t : SM \rightarrow SM$ be the geodesic flow. The natural projection of the tangent bundle will be denoted by $\pi : TM \rightarrow M$. We will write elements of the tangent bundle either as sole vectors v (which is in $T_x M$) or as pairs (x, v) , whichever is more convenient.

The boundary of the sphere bundle is

$$\partial SM = \{v \in SM : \pi(v) \in \partial M\}.$$

We identify the inward-pointing part of this boundary,

$$\partial_{\text{in}} SM = \{v \in \partial SM : \langle v, \nu \rangle_v > 0\},$$

where ν is the inward-pointing normal vector field and $\langle v, \nu \rangle_v := g_{ij}(v)v^i v^j$. Similarly, we define the outward-pointing part

$$\partial_{\text{out}} SM = \{v \in \partial SM : \langle v, \nu \rangle_v < 0\}.$$

Definition 3. Let (M, F) be a Finsler manifold with boundary. For each $t > 0$ we define a relation R_t on $\partial_{\text{in}} SM$ so that $v R_t w$ if and only if there exist two numbers $t_1, t_2 > 0$ for which $t_1 + t_2 = t$ and $\pi(\phi_{t_1}(v)) = \pi(\phi_{t_2}(w))$. We call this relation the broken scattering relation.

Our main result is the following theorem stating that the broken scattering relation, that is, the lengths of the broken geodesics, determine uniquely the isometry type of a Finsler manifold.

Theorem 4. Let $(M_i, F_i), i \in \{1, 2\}$, be two compact Finsler manifolds of dimension larger or equal to 3, with boundary. We assume the following:

- (i) Both Finsler functions F_1 and F_2 are reversible.
- (ii) The manifolds $(M_i, F_i), i \in \{1, 2\}$, have strictly convex foliations in the sense of [Definition 1](#).
- (iii) There are diffeomorphisms $\xi : \partial M_1 \rightarrow \partial M_2$ and $\Xi : \partial TM_1 \rightarrow \partial TM_2$ that are compatible in the sense of [Definition 2](#).
- (iv) $F_1 = F_2 \circ \Xi$ on ∂TM_1 .
- (v) For any two vectors $v, w \in \partial_{\text{in}} SM_1$ and $t > 0$, we have $v R_t^{(1)} w$ if and only if $\Xi(v) R_t^{(2)} \Xi(w)$, where $R_t^{(i)}$ is the broken scattering relation of [Definition 3](#) on (M_i, F_i) .

Then there is a diffeomorphism $\phi : M_1 \rightarrow M_2$ that is an isometry in the sense of $F_1 = F_2 \circ d\phi$, which satisfies $\phi|_{\partial M_1} = \xi$ and $d\phi|_{\partial TM_1} = \Xi$.

Remark 5. We note that to prove [Theorem 4](#) we will actually only need that (M_1, F_1) has strictly convex foliation and (M_2, F_2) has strictly convex boundary.

The requirement for the dimension $n \in \mathbb{N}$ of M_i being larger than 2 is a technical one. This is, however, a vital component of the proof of [Lemma 19](#). In the proof we need to assume that the dimension of a trace of geodesic, which is 1, is strictly smaller than the dimension of the boundary ∂M_i , which is $n - 1$.

There are several well-known examples of compact Riemannian manifolds satisfying a convex foliation condition of [Definition 1](#). One example that is particularly interesting to us is the closed unit ball $\bar{B} \subset \mathbb{R}^n$ with a radial metric $g(x) = c^{-2}(r)e$, where $r = |x| \in [0, 1]$. Here c is a strictly positive smooth function on $[0, 1]$ satisfying the Herglotz condition

$$\frac{d}{dr} \left(\frac{r}{c(r)} \right) > 0$$

and $c'(0) = 0$. The condition does not forbid the existence of conjugate points but there are no trapped geodesics [[Herglotz 1905](#)].

Consider the smooth function $f(x) = |x|^2 - 1$ on \bar{B} . It is well known that the radial Riemannian metric $g(x) = c^{-2}(|x|)e$ satisfies the Herglotz condition if and only if the function f satisfies the condition of [Definition 1](#). Due to convexity, in terms of [Definition 1](#), being an open property we have that the function f is a strictly convex foliation of a Finsler manifold (\bar{B}, F) for any Finsler function F on \bar{B} that is close enough to g in the C^2 -topology.

Rotation-symmetric Riemannian manifolds satisfying the Herglotz condition can have conjugate points. Therefore some small Finslerian variations of such manifolds, at least those that preserve rotation symmetry, also have conjugate points. Thus our conditions do allow for conjugate points on non-Riemannian manifolds.

1.1. Comparison of inverse problems for Riemannian and Finsler manifolds and the Riemannian counterpart of [Theorem 4](#). Several inverse problems studied in the literature have a different nature in Riemannian and Finslerian settings. For example, recall that a Riemannian surface is simple if it is simply connected, its geodesics have no conjugate points and ∂M is strictly convex. For the travel time inverse problem, discussed in detail below, it is shown in [[Pestov and Uhlmann 2005](#)] that the isometry type of a simple Riemannian manifold is uniquely determined by the distances $d(x, y)$ of the boundary points $x, y \in \partial M$, but there are counterexamples for this problem for simple Finsler manifolds [[Ivanov 2013](#)]. Another example of the different nature of Riemannian and Finslerian settings is that the blow-up maps have been used to construct nonsmooth counterexamples [[Greenleaf et al. 2003; 2009](#)] for inverse problems for the Laplace–Beltrami operator, corresponding to a Riemannian metric. However, these maps cannot be used for elastic equations, corresponding to a Finsler structure, as elastic media has to satisfy physical symmetries [[Sklan et al. 2018](#)].

The Riemannian counterpart of [Theorem 4](#), presented in [[Kurylev et al. 2010](#)], needs fewer assumptions. This is due to a certain rigidity of Riemannian geometry. Heuristically, in Finsler geometry the metrics in different directions at the same point are independent. This implies that in order to reconstruct a Finsler manifold in full, one needs access to the entire tangent bundle. To prove [Theorem 4](#) the first step is to

show that the broken scattering relation determines the collection of distance functions at the boundary (see [Proposition 7](#)). However, these distances only give access for a certain open set of directions.

The important feature of a Finsler function arising from a Riemannian metric tensor is that it is real-analytic on every punctured tangent space. If one makes this additional assumption, then the distance functions at the boundary determine the geometry at all points in all directions as proven in [\[Hoop et al. 2019\]](#). By [Lemma 9](#) the distance function determines the Finsler function in directions where the geodesic is minimal to its endpoint on the boundary. We call this the good subset of the tangent bundle. These good directions exist on every tangent space, which is why analyticity gives access to the entire bundle.

The main result of [\[Hoop et al. 2019\]](#) also asserts that the collection of the distance functions does not determine the Finsler function outside the good set of the tangent bundle. Therefore on a general Finsler manifold one needs to study also long geodesics, and we have to go beyond the scope of [\[Kurylev et al. 2010\]](#). Reversibility and strict convexity give access to all directions near the boundary. To go further into the manifold, we use the foliation: every point in the interior is close to the boundary when one goes deep enough in the foliation.

1.2. Elasticity and Finsler geometry. One can define elastic geometry in terms of distance: the distance between two points can be declared to be the shortest time it takes for the support of a solution to the elastic wave equation, originally supported at one point, to reach the other point.

A more tractable description is obtained by studying the propagation of singularities of the elastic wave equation. There is a concrete way to pass from the stiffness tensor to a Minkowski norm on each tangent space, and this is described in [\[Hoop et al. 2019\]](#). Microlocal analysis indicates [\[Duistermaat 1996; Greenleaf and Uhlmann 1993\]](#) that singularities follow the geodesic flow on the cosphere bundle of a Finsler metric, and cospheres are known as slowness surfaces in the physical literature. Elastic waves have three different polarizations in three spatial dimensions. The singularities used in the derivation of the Finsler geometry correspond to the fastest polarization known as quasipressure or qP.

In seismology the commonly used preliminary reference earth model [\[Dziewonski and Anderson 1981\]](#) is spherically symmetric and satisfies a foliation condition, the Herglotz condition, to great accuracy. A typical inverse problem in elasticity is to reconstruct some elastic properties from boundary data [\[Bal et al. 2015; Bao et al. 2018; Barceló et al. 2018; Beretta et al. 2014; Hu et al. 2014; Liu and Xiao 2017; Mazzucato and Rachele 2007; Nakamura and Uhlmann 2003\]](#). Given this geometric point of view, the task is to find the Finsler geometry corresponding to the parameters.

It was shown in [\[Hoop et al. 2019\]](#) that Finsler metrics arising from elasticity are always reversible and fiberwise real-analytic. The following result is an adaptation of [Theorem 4](#) in the elastic setting:

Theorem 6. *Let (M_i, F_i) , $i \in \{1, 2\}$, be two compact Finsler manifolds of dimension larger or equal to 3, with strictly convex boundaries. We assume the following:*

- (i) *Both Finsler functions F_1 and F_2 are reversible.*
- (ii) *The manifolds (M_i, F_i) , $i \in \{1, 2\}$, are fiberwise real-analytic. That is, for every $x \in M_i$ the function*

$$F_i(x, \cdot) : T_x M_i \rightarrow \mathbb{R}$$

is real-analytic.

- (iii) *There are diffeomorphisms $\xi : \partial M_1 \rightarrow \partial M_2$ and $\Xi : \partial TM_1 \rightarrow \partial TM_2$ that are compatible in the sense of [Definition 2](#).*
- (iv) $F_1 = F_2 \circ \Xi$ on ∂TM_1 .
- (v) *For any two vectors $v, w \in \partial_{\text{in}} SM_1$ and $t > 0$ we have $v R_t^{(1)} w$ if and only if $\Xi(v) R_t^{(2)} \Xi(w)$, where $R_t^{(i)}$ is the broken scattering relation of [Definition 3](#) on (M_i, F_i) .*

Then there is a diffeomorphism $\phi : M_1 \rightarrow M_2$ that is an isometry in the sense of $F_1 = F_2 \circ d\phi$, which satisfies $\phi|_{\partial M_1} = \xi$ and $d\phi|_{\partial TM_1} = \Xi$.

In the elastic setting no foliation condition is needed; it can be replaced with fiberwise analyticity. But we do point out that with the foliation condition we have direct access to the Finsler function on all of the tangent bundle, whereas without the foliation condition we have to resort to analytic continuation which may be problematic for applications. If our methods are used outside Hookean elasticity, the analyticity property and reversibility may no longer be available.

1.3. Related problems. Let us consider a compact Riemannian manifold (M, g) with boundary. The classical boundary rigidity problem is the following: We assume that we are given the distances $d(x, y)$, through M , of all boundary points $x, y \in \partial M$. Can we determine the isometry type of the manifold (M, g) ? Michel [[1981](#); [1994](#)] observed that in the case of simple manifolds these distance functions also determine the values of the geodesic flow at the boundary, this is the scattering relation or lens relation:

$$L = \{(v, w, t) \in \partial_{\text{in}} SM \times \partial_{\text{out}} SM \times \mathbb{R} : \phi_t(v) = w \text{ for some } t \leq 0\}.$$

Thus L carries the information of when and where and in which direction a geodesic, sent from the boundary, hits the boundary again.

The natural conjecture is that for simple manifolds the scattering relation determines the isometry type of the manifold; see [[Burago and Ivanov 2010](#); [Croke 1991](#); [Gromov 1983](#); [Lassas et al. 2003](#); [Michel 1981](#); [Muhometov 1981](#); [Pestov and Uhlmann 2005](#); [Stefanov and Uhlmann 2005](#)]. If the manifold is trapping one cannot determine the metric up to isometry if only the scattering relation is known [[Croke and Kleiner 1994](#)].

On the other hand if a Riemannian manifold admits a suitable convex foliation condition, then a local version of the lens rigidity problem, studied in [[Stefanov et al. 2016](#); [2021](#); [Uhlmann and Vasy 2016](#)], implies the global boundary rigidity result. See for instance [[Paternain et al. 2019](#), Section 2] for a survey of different types of foliation conditions and geometric properties that imply their existence.

Microlocal analysis connects singularities of solutions and solution operators to (hyperbolic) partial differential equations (PDEs) to geometry. The PDE related to our problem is the elastic wave equation. Its principal symbol is a matrix whose largest eigenvalue is directly related to the Finsler metric [[Hoop et al. 2019](#), Section 2]. Single scattering can be modeled by introducing, in the elastic wave equation, a right-hand side representing a contrast source. The coefficients in this source, identified with the contrast in stiffness tensor, have a nonempty wavefront set. In the inverse problem studied in this paper, this wavefront set is assumed to be dense on the cosphere bundle associated with M . Thus in the case of very heterogeneous media with many scattering points inside the manifold, one can obtain further

information by looking at the propagation of singularities of waves going through the manifold. This is the broken scattering relation. Kurylev, Lassas and Uhlmann [Kurylev et al. 2010] showed that this relation determines the Riemannian manifold (M, g) , up to an isometry. They reduce the problem to the setup of [Kurylev 1997; Katchalov et al. 2001], where it is shown that the collection of distance functions at the boundary determines the isometry class. In a sense they turn a data given by boundary sources to one given by interior point sources. For inverse problems related to interior point sources and Riemannian wave equation, see for instance [de Hoop and Saksala 2019; Ivanov 2020; Lassas and Saksala 2019; Lassas et al. 2018; Oksanen 2011], and [Hoop et al. 2019] for an interior source problem in Finsler geometry.

2. Outline of the proof

We prove [Theorem 4](#) in this section. The proofs of the key lemmas are postponed to subsequent sections. The rough plan of the proof is as follows:

- The first step is to verify that the broken scattering relation determines the boundary distance functions $\{d(x, \cdot) : \partial M \rightarrow \mathbb{R} \mid x \in \text{int } M\}$. Then we show that the boundary distance function determines the topological and smooth structures. This is based on earlier work [Hoop et al. 2019].
- By the previous result the boundary distance function determines F on the part of TM from where the geodesic flow reaches the boundary in a sufficiently short time. This part is known as the “good set” $G \subset TM \setminus 0$.
- If a point $x \in M$ is sufficiently close to the strictly convex boundary, then more than half of the directions on $T_x M$ belong to G , so F is determined there.
- By reversibility, F is determined on all of $T_x M$. This implies that we have found the Finsler geometry in all directions in a neighborhood of the boundary. Using the foliation, we write the neighborhood as $f^{-1}([0, \epsilon])$ for some $\epsilon > 0$.
- We may then ask how far in the foliation the metric is uniquely determined. By the previous argument it holds at least for a little bit. If it only holds up to some $s < S$, then we may reiterate the argument on the smaller manifold $f^{-1}([s, S])$. To do so, we must propagate the data from the original boundary inward to the new one. Thus the uniqueness extends beyond the alleged limit s , proving uniqueness on the whole manifold.

We first give the lemmas and definitions needed to make the proof precise, then finish the proof, and finally complete the proof by proving the lemmas.

2.1. Auxiliary results. The boundary distance function of a point $x \in M$ is the function $r_x : \partial M \rightarrow \mathbb{R}$ defined by $r_x(y) = d(x, y)$. Notice that we assumed the Finsler metrics to be reversible, so it does not matter which direction we measure the distance in. In the cases where we are considering multiple manifolds, the boundary distance function of $x \in M_i$ is denoted by $r_x^{(i)}$.

Proposition 7 (proven in [Section 5](#)). *Let (M_i, F_i) , $i = 1, 2$, be two compact reversible Finsler manifolds, with strictly convex boundaries, whose broken scattering relations agree in the sense of conditions (iii)–(v)*

in [Theorem 4](#). Then the boundary distance functions agree in the sense that

$$\{r_x^{(1)} : x \in \text{int } M_1\} = \{r_y^{(2)} \circ \xi : y \in \text{int } M_2\}. \tag{1}$$

Remark 8. We note that in (1) we are given nonindexed sets. That is, for a function $r_x : \partial M \rightarrow \mathbb{R}$ we know whether it is included in the set (1), but we do not know the indexing point $x \in M$. The map $\xi : \partial M_1 \rightarrow \partial M_2$ is the same as in [Theorem 4](#).

We say that a direction $v \in TM \setminus 0$ is minimizing if the maximal geodesic starting at v reaches ∂M in finite time and is a shortest curve joining its endpoints. We denote the set of minimizing directions by G (for “good”)—obviously with $G^{(i)} \subset TM_i \setminus 0$. The fibers are denoted by $G_x^{(i)}$.

Lemma 9 [[Hoop et al. 2019](#), Theorem 1.3]. *Let (M_1, F_1) and (M_2, F_2) be two compact Finsler manifolds with boundary. Suppose there is a diffeomorphism $\xi : \partial M_1 \rightarrow \partial M_2$ so that (1) holds. Then there is a diffeomorphism $\phi : M_1 \rightarrow M_2$ so that $\phi|_{\partial M_1} = \xi$. In addition, $F_1(v) = F_2(d\phi(v))$ for all $v \in G^{(1)}$.*

For a set $A \subset T_x M$ we denote by $-A$ the reflection and by $\sigma(A) = A \cup -A$ the symmetrization. We also use the same notation on the whole bundle, applied fiberwise.

Lemma 10 (proven in [Section 3](#)). *If a Finsler manifold (M, F) has a strictly convex boundary, then the set*

$$U = \{x \in M : \sigma(G_x) = T_x M \setminus 0\}$$

is a neighborhood of the boundary $\partial M \subset M$.

Lemma 11 (proven in [Section 4](#)). *Let F_1 and F_2 be two Finsler functions on a compact smooth manifold M with boundary. Suppose (M, F_1) has a strictly convex foliation with a function $f : M \rightarrow [0, S]$. Let $s \in (0, S)$ and set $\widehat{M} = f^{-1}([s, S])$. Assume the following:*

- *The two metrics coincide below s in the sense that $F_1 = F_2$ on $T(M \setminus \widehat{M})$.*
- *For any two $v, w \in \partial_{\text{in}} SM$ (these bundles coincide for the two metrics) and $t > 0$ we have $vR_t^{(1)} w$ if and only if $vR_t^{(2)} w$, where $R_t^{(i)}$ is the broken scattering relation of (M, F_i) .*

Then the scattering relations $\widehat{R}_t^{(i)}$ on (\widehat{M}, F_i) coincide in the sense that the assumptions of [Theorem 4](#) are valid with ξ and Ξ being the identity maps. Most importantly, $F_1 = F_2$ on $\partial T\widehat{M}$ and for all $v, w \in \partial_{\text{in}} S\widehat{M}$ we have $v\widehat{R}_t^{(1)} w$ if and only if $v\widehat{R}_t^{(2)} w$.

Finally we recall the second main result of [[Hoop et al. 2019](#)]:

Lemma 12 [[Hoop et al. 2019](#), Theorem 1.5]. *Let (M_1, F_1) and (M_2, F_2) be two compact Finsler manifolds with boundary. Suppose there is a diffeomorphism $\xi : \partial M_1 \rightarrow \partial M_2$ so that (1) holds. If Finsler functions F_1 and F_2 are fiberwise real-analytic, then there exists a Finslerian isometry $\Psi : (M_1, F_1) \rightarrow (M_2, F_2)$ so that $\Psi|_{\partial M_1} = \phi$.*

2.2. Proofs of the theorems. Now we are ready to present the detailed proofs of [Theorems 4](#) and [6](#) respectively. We use the notation introduced in the previous subsection.

Proof of Theorem 4. It follows from Proposition 7 that the boundary distance functions of the two manifolds agree up to identifying the boundaries with ξ . By Lemma 9 there is a diffeomorphism $\phi : M_1 \rightarrow M_2$, and we use this diffeomorphism to identify the two manifolds as $M = M_1 = M_2$. This manifold inherits the foliation from (M_1, F_1) .

This smooth manifold M has two Finsler functions F_1 and F_2 , and the goal is to show that they are equal. Lemma 9 shows that $F_1 = F_2$ in the good set $G := G^{(1)} \subset TM \setminus 0$. By reversibility we conclude that $F_1 = F_2$ in $\sigma(G)$. Lemma 10 guarantees that $\sigma(G)$ contains the punctured tangent bundle of some neighborhood of ∂M . Therefore there is $\epsilon > 0$ so that $F_1 = F_2$ in tangent bundle $T(f^{-1}([0, \epsilon]))$.

Now define

$$I = \{h \in [0, S] : F_1(x, v) = F_2(x, v) \text{ whenever } f(x) \leq h \text{ and } v \in T_x M \setminus 0\}.$$

We showed that $[0, \epsilon) \subset I$, and it is clear that I is an interval. By continuity of the two Finsler functions, it is a closed interval and therefore $I = [0, s]$ for some $s \in (0, S]$. If $s = S$, the two Finsler functions coincide on the whole tangent bundle and the proof is complete.

Suppose then that $s \in (0, S)$. We are now in the setting of Lemma 11. The two metrics coincide in the strip $f^{-1}([0, s])$, so the data may be propagated through it. We have now two metrics on the shrunken manifold $\widehat{M} = f^{-1}([s, S])$ and their broken scattering relations coincide by Lemma 11.

Let us denote the good set of (\widehat{M}, F_i) by $\widehat{G}^{(i)}$. Repeating the argument obtained above, we find that there is a diffeomorphism $\eta : \widehat{M} \rightarrow \widehat{M}$ so that $F_1 = \eta^* F_2$ on $\widehat{G}^{(1)}$, which again has full fibers when the base point is in some neighborhood of the boundary. It is straightforward to check that if $(x, v) \in G$ and $x \in \widehat{M}$, then $(x, v) \in \widehat{G}^{(1)}$. We also note that for any $x \in \widehat{M}$ the intersection $T_x \widehat{M} \cap G^{\text{int}}$ is nonempty, since any small perturbation of an initial direction of the geodesic connecting x to a closest boundary point in ∂M , is contained in G . Take $(x, v) \in G^{\text{int}}$ and let γ be the forward-maximal geodesic with respect to F_1 with such initial data. Due to [Hoop et al. 2019, Section 3.3] it follows that the lift of γ is contained in G^{int} . Since the geodesic coefficients $G^i(w)$, $i \in \{1, \dots, n\}$, $w \in TM$ (see for instance [Shen 2001, Formula (5.7)] for the definition), of F_1 and F_2 agree on G^{int} , it follows that γ is also a geodesic with respect to F_2 .

But as F_1 and $\eta^* F_2$ agree in a neighborhood of the lift of γ , we have that the F_2 -geodesic γ starts at $\eta(x)$. Thus $\eta(x) = x$ and one could choose any starting point $x \in \widehat{M}$, and so $\eta = \text{id}$. Therefore $F_1(x, v) = F_2(x, v)$ for $x \in \widehat{M}$ close enough to $\partial \widehat{M}$ and all $v \in T_x M$. This means that the two metrics coincide in $f^{-1}([0, s + \epsilon])$ for some $\epsilon > 0$, contradicting the maximality of s .

Finally we verify that $d\phi$ and Ξ coincide on ∂TM_1 . First we note that Definition 2, Proposition 7 and Lemma 9 imply

$$d\phi|_{T \partial M_1} = d\xi = \Xi|_{T \partial M_1}. \tag{2}$$

We recall that Ξ and $d\phi$ are linear on the fibers, and therefore to conclude the proof it suffices to show that Ξ preserves the inward-pointing unit normal vector field to the boundary. By taking the directional differential d_v of condition (iv) of Theorem 4, and using equation (2), we get $\langle d_v(F_2)(\Xi v), d\xi w \rangle = 0$, $w \in T \partial M_1$.

Since $\xi : \partial M_1 \rightarrow \partial M_2$ is a diffeomorphism we have shown that the Legendre transform of Ξv is conormal to $T \partial M_2$. This implies that Ξv is normal to the boundary ∂M_2 . Due to condition (v) of

Theorem 4 it holds that Ξ maps $\partial_{\text{in}}SM_1$ onto $\partial_{\text{in}}SM_2$. Therefore we have verified $\Xi\nu$ is the inward-pointing unit normal to ∂M_2 . \square

Proof of Theorem 6. **Theorem 6** is a direct consequence of **Proposition 7** and **Lemma 12**. \square

3. Short and long geodesics

In this section we analyze short geodesics near the boundary to prove **Lemma 10**. To do so, we first make an observation concerning the lengths of geodesics of arbitrary — even infinite — length.

For any Finsler manifold (M, F) with boundary we define the exit time function $\tau_{\text{exit}} : SM \rightarrow [0, \infty]$ so that $\tau_{\text{exit}}(v)$ is the first time $t > 0$ for which $\pi(\phi_t(v)) \in \partial M$. If there is no such time, the value is taken to be infinity. If v is based at a boundary point and points tangentially or outward, we set $\tau_{\text{exit}}(v) = 0$.

Lemma 13. *On any metrically complete Finsler manifold M with strictly convex boundary the exit time function $\tau_{\text{exit}} : SM \rightarrow [0, \infty]$ is continuous.*

Without assuming the metric completeness, continuity holds where $\tau_{\text{exit}} < \infty$, including a neighborhood of $\overline{\partial_{\text{out}}SM}$.

Proof. We will check continuity separately at different kinds of points $v \in SM$, depending on the value $\tau_{\text{exit}}(v)$.

If $\tau_{\text{exit}}(v) = 0$, then v is based on the boundary and points tangentially or outwards. (That is, $v \in \overline{\partial_{\text{out}}SM}$.) By the assumption of strict convexity, any geodesic starting near v is short due to [**Stefanov and Uhlmann 2018**, Section 8.1], establishing continuity of τ_{exit} at v .

If $\tau_{\text{exit}}(v) \in (0, \infty)$, then the geodesic γ_v starting at v reaches ∂M in finite time. By strict convexity $\dot{\gamma}_v(\tau_{\text{exit}}(v))$ is transverse to ∂M , and it follows from the implicit function theorem that τ_{exit} is in fact smooth in a neighborhood of v .

If $\tau_{\text{exit}}(v) = \infty$, then the geodesic γ_v is trapped. Continuity at v can only fail if there is a sequence of vectors $v_k \in SM$ so that $v_k \rightarrow v$ as $k \rightarrow \infty$ and $\tau_{\text{exit}}(v_k) \leq T$ for some $T \in (0, \infty)$. There is some $r > 0$ so that for all $k \in \mathbb{N}$ we have $d(\pi(v_k), \pi(v)) < r$. All the forward-maximal geodesics γ_{v_k} are thus contained in the metric closed ball $K = \overline{B}(\pi(v), r + T)$. By completeness of M the set K is compact. Up to extracting a subsequence, the endpoints $\gamma_{v_k}(\tau_{\text{exit}}(v_k)) \in \partial M \cap K$ converge to a point $z \in \partial M$ and $\tau_{\text{exit}}(v_k)$ converges to $T' \leq T$. By continuity of the geodesic flow, we have that $\phi_{T'}(v) = \lim_{k \rightarrow \infty} \phi_{\tau_{\text{exit}}(v_k)}(v_k) \in S_z M$. This means that γ_v meets the boundary at time $T' < \infty$, which is a contradiction.

We conclude that τ_{exit} is continuous at all points of SM . Completeness was only needed to prove continuity where $\tau_{\text{exit}} = \infty$, and that cannot happen near $\overline{\partial_{\text{out}}SM}$ when ∂M is strictly convex. \square

Proof of Lemma 10. If the manifold is not compact, we can restrict the analysis to a bounded neighborhood of a given boundary point and do everything on a compact submanifold. On a compact Finsler manifold there is some $\epsilon > 0$ so that any geodesic with length less than ϵ is minimizing.

Consider any point $p \in \partial M$, and let ν_p be the inward unit normal at p and γ_p the geodesic starting in the direction of ν_p . For any $t > 0$ small enough so that the geodesic γ_p up to time t exists we define the map $P_t^p : S_p M \rightarrow S_{\gamma_p(t)} M$ by parallel transport.

Fix some $T > 0$ small enough. We define a function $Q : [0, T) \times \partial SM \rightarrow SM$ so that $Q(t, v) = P_t^{\pi(v)}(v)$. This function is continuous, so by Lemma 13 the composed function $\tau_{\text{exit}} \circ Q : [0, T) \times \partial SM \rightarrow [0, \infty]$ is also continuous. (Continuity will only be needed near directions where $\tau_{\text{exit}} = 0$.) This composed function vanishes on $\{0\} \times \overline{\partial_{\text{out}} SM}$, so there is an open neighborhood $V \subset [0, T) \times SM$ of $\{0\} \times \partial_{\text{out}} SM$ so that $\tau_{\text{exit}} \circ Q|_V < \epsilon$. Since all geodesics shorter than ϵ are minimizing, we have $Q(V) \subset G$.

Any boundary point $p \in \partial M$ has a neighborhood $W \subset \partial M$ and two numbers $a, h > 0$ so that $[0, h) \times V_p \subset V$ with

$$V_p := \{v \in SM : \pi(v) \in W \text{ and } \langle v, v \rangle_v < a\}.$$

Note that V_p contains a larger portion of $S_p M$ than the Southern hemisphere. We have $Q([0, h) \times V_p) \subset G$ and

$$Q([0, h) \times \sigma(V_p)) = \sigma(Q([0, h) \times V_p)) \subset \sigma(G).$$

But we have $\sigma(V_p) = \{v \in SM : \pi(v) \in W\}$, so that the set $U' = \{\gamma_p(t) : t \in [0, h) \text{ and } p \in W\}$ is a subset of the set U defined in the claim of the lemma. Now U' is a neighborhood of the arbitrary boundary point p , so we have established that U does indeed contain a neighborhood of the boundary. \square

4. Propagation of data through a layer

In order to prove Lemma 11 to propagate data from ∂M to $\partial \widehat{M}$, we first observe that a foliated manifold enjoys a certain weak nontrapping property. However, these manifolds can have trapped geodesics as shown via examples given in the discussion after Definition 1.

Lemma 14. *Let (M, F) be a compact Finsler manifold which is foliated by a smooth function $f : M \rightarrow \mathbb{R}$ in the sense of Definition 1. If a maximal geodesic γ satisfies $\partial_t(f(\gamma(t))) < 0$ at $t = 0$, then γ reaches ∂M in finite time.*

Proof. Consider the function $h(t) = f(\gamma(t))$, defined over some maximal interval $[0, T]$ or $[0, \infty)$. The goal is to show that h obtains the value zero in finite time. Given that $h'(0) < 0$ and h is smooth, this can only fail if one of the following happen:

- (i) The derivative $h'(t)$ vanishes for some t .
- (ii) The function has a limit: h is strictly decreasing and $\lim_{t \rightarrow \infty} h(t) = H \in [0, \infty)$.

Let us first exclude case (i). If h' vanishes somewhere, there is a smallest $t > 0$ for which $h'(t) = 0$. We have that $h' < 0$ on $[0, t)$. The geodesic γ is tangent to the level set $f^{-1}(h(t))$, so by Definition 1 we have $h''(t) < 0$. This implies that for some $\epsilon > 0$ we have $h'(t - \epsilon) > 0$, which is a contradiction.

Let us then move to case (ii). The curve γ now approaches the surface $\Sigma := f^{-1}(H)$. By monotonicity there is an increasing sequence of times $t_k \rightarrow \infty$ so that $h'(t_k) \rightarrow 0$. Upon extracting a subsequence, there exists a point $x \in \Sigma$ so that $\gamma(t_k) \rightarrow x$ and $\dot{\gamma}(t_k)$ converges to some $v \in S_x M$. As $h'(t_k) \rightarrow 0$, it follows that v is tangent to Σ .

Let $\tilde{\gamma}$ be the geodesic starting with initial conditions (x, v) . We set $\tilde{h} = f \circ \tilde{\gamma}$. Since v is tangent to Σ , we have $\tilde{h}'(0) = 0$, so Definition 1 implies $\tilde{h}''(0) < 0$. By the smoothness of f and the time additive

property of geodesic flow we have

$$\lim_{k \rightarrow \infty} h''(t_k + s) = \tilde{h}''(s)$$

for any $s \in \mathbb{R}$.

Because $\tilde{h}''(0) < 0$, there are $\epsilon > 0$ and $\delta > 0$ so that $h''(t_k + s) \leq -\epsilon$ for all $s \in [-\delta, \delta]$ when k is large enough. This together with $h' \leq 0$ (which is due to monotonicity) and the fundamental theorem of calculus gives

$$h(t_k + \delta) = h(t_k - \delta) + 2\delta h'(t_k - \delta) + \int_{-\delta}^{\delta} \int_{-\delta}^s h''(t_k + r) dr ds \leq h(t_k - \delta) - 2\epsilon\delta^2$$

for large enough k . As $-2\epsilon\delta^2$ is independent of k and h is monotonous, it follows that h cannot converge. This is a contradiction.

Therefore there is indeed some finite t so that $h(t) = 0$. □

Proof of Lemma 11. Equality of the Finsler functions on $\partial\widehat{M}$ to all directions on the bundle — which amounts to ξ and Ξ being identities — follows from the assumed equality of the Finsler functions on the strip $f^{-1}([0, s])$ and continuity.

For any $v \in SM$, we denote by γ_v the unique geodesic with $\dot{\gamma}_v(0) = v$. Take any two $v, w \in \partial_{\text{in}}S\widehat{M}$ and $t > 0$.

By Lemma 14 there is $a > 0$ so that the geodesic segment $\gamma_v|_{[-a, 0]} \subset f^{-1}([0, s])$ connects a point on ∂M to the point $\pi(v) \in \partial\widehat{M}$. Because the two Finsler metrics agree in $f^{-1}([0, s])$, this same curve is a geodesic in both geometries. By strict convexity, this geodesic meets ∂M transversely, so that $v' := \dot{\gamma}_v(-a)$ belongs to $\partial_{\text{in}}SM$. Similarly, there are $b > 0$ and $w' \in \partial_{\text{in}}SM$ corresponding to w .

Because the two geodesic flows agree from v' to v and from w' to w , we have that $v\widehat{R}_t^{(i)}w$ if and only if $v'R_{t+a+b}^{(i)}w'$ for both $i \in \{1, 2\}$. The broken scattering relations $R_{t+a+b}^{(i)}$ agree on ∂M , and so the broken scattering relations $\widehat{R}_t^{(i)}$ agree on $\partial\widehat{M}$. □

5. From broken scattering relation to boundary distance functions

In this section we will prove Proposition 7. We start with considering only one Finsler manifold (M, F) with strictly convex boundary whose broken scattering relation is known.

5.1. Critical distance functions. We begin with studying classical critical distance functions. The first one is the cut distance function $\tau_{\text{cut}} : SM \rightarrow (0, \infty]$ given by

$$\tau_{\text{cut}}(x, v) := \sup\{t > 0 : \gamma_{x,v}(t) \text{ exists and } d(x, \gamma_{x,v}(t)) = t\}.$$

Since the boundary of the M is strictly convex it holds that any distance-minimizing curve is a geodesic. Therefore function τ_{cut} is well-defined and moreover it is continuous.

Next we formulate two auxiliary lemmas related to τ_{cut} .

Lemma 15. *For any $(x, v) \in \partial_{\text{in}}SM$ and $s_2 > s_1 \geq 0$ which satisfy*

$$\gamma_{x,v}(s_1) = \gamma_{x,v}(s_2), \quad \text{we have } s_1 + s_2 > 2\tau_{\text{cut}}(x, v).$$

Proof. Since F is reversible this claim can be proven like [Kurylev et al. 2010, Lemma 2.1]. □

Lemma 16. *Let $z \in \partial M$, $t \in (0, \tau_{\text{cut}}(z, \nu(z)))$ and $U \subset TM$ be a neighborhood of $\nu(z)$ that is diffeomorphic to some open set of $\mathbb{R}^{2n-1} \times [0, \infty)$. For any $\epsilon > 0$ we can choose $\delta = \delta(z, t, \epsilon) > 0$ such that the following holds: If $v_i \in U$, $i \in \{1, 2\}$, satisfy*

$$v_1 R_{2t} v_2 \quad \text{and} \quad \|v_i - \nu(z)\|_e < \delta,$$

then there exist $t_1, t_2 > 0$ so that

$$\gamma_{v_1}(t_1) = \gamma_{v_2}(t_2), \quad t_1 + t_2 = 2t, \quad \text{and} \quad |t_i - t| < \epsilon.$$

Here $\|\cdot\|_e$ is the Euclidean norm on U .

Proof. The proof is analogous to the proof of [Kurylev et al. 2010, Lemma 2.2]. □

The second critical distance function is the boundary cut distance function

$$\tau_{\partial M}(z) := \sup\{t > 0 : d(z, \gamma_{z, \nu(z)}(t)) = t = d(\partial M, \gamma_{z, \nu(z)}(t))\}, \quad z \in \partial M.$$

These two critical distance functions satisfy the following:

Lemma 17 [Hoop et al. 2019, Lemma 3.8]. *For any $z \in \partial M$ it holds that*

$$\tau_{\text{cut}}(z, \nu(z)) > \tau_{\partial M}(z).$$

5.2. Family of focusing directions. Let (M, F) be a compact Finsler manifold of dimension 3 or higher with reversible Finsler function F and strictly convex boundary. Let $x \in M^{\text{int}}$ and $v \in S_x M$ be a direction such that the corresponding geodesic γ_v is the distance minimizer from x to $z_x \in \partial M$, a closest boundary point to x . In the proof of Lemma 13 we showed that the exit time function τ_{exit} is smooth in some neighborhood $V \subset S_x M$ of v , and, moreover, the implicit function theorem implies that the map

$$H : V \rightarrow \partial M, \quad \eta \mapsto \pi(\phi_{\tau_{\text{exit}}(\eta)}(\eta)),$$

is a diffeomorphism to some neighborhood $U \subset \partial M$ of z_x . Using this identification we define a smooth inward-pointing unit length vector field on U

$$V(z) := -\phi_{\tau_{\text{exit}}(\eta)}(\eta), \quad \eta := H^{-1}(z).$$

This vector field satisfies

$$V(z_1) R_{T(z_1, z_2)} V(z_2) \quad \text{and} \quad V(z_x) = \nu(z_x),$$

where

$$T(z_1, z_2) := \tau_{\text{exit}}(H^{-1}(z_1)) + \tau_{\text{exit}}(H^{-1}(z_2)).$$

Moreover, the function

$$t : U \rightarrow \mathbb{R}, \quad t(z) := \frac{1}{2} T(z, z) = \tau_{\text{exit}}(H^{-1}(z))$$

is smooth and its differential vanishes at z_x which is a closest boundary point to x . Thus the geodesics given by initial conditions $(z, V(z))$, $z \in U$, focus at the common interior point x at time $t(z)$.

We change the point of view and set the following definition:

Definition 18. Let $z_0 \in \partial M$ and $t_0 \in (0, \tau_{\text{exit}}(\nu(z_0)))$. We say that a collection

$$F(z_0, t_0) := \{U, V(\cdot), t(\cdot)\},$$

where $U \subset \partial M$ is a neighborhood of z_0 , $V : U \rightarrow \partial_{\text{in}}SM$ is a smooth vector field and $t : U \rightarrow (0, \infty)$ is a smooth function, is called a family of focusing directions around (z_0, t_0) if

$$V(z_1)R_{t(z_1)+t(z_2)}V(z_2) \quad \text{for all } z_1, z_2 \in U, \tag{3}$$

$$V(z_0) = \nu(z_0), \quad t(z_0) = t_0 \quad \text{and} \quad dt(z)|_{z=z_0} = 0. \tag{4}$$

We note that the broken scattering relation determine all the families of focusing directions, but it is possible that not all of them focus in the sense of

$$\pi(\phi_{t(z)}(V(z))) = \pi(\phi_{t_0}(\nu(z_0))) \quad \text{for all } z \in U. \tag{5}$$

Next we give the following result that guarantees the actual focusing if the focusing time t_0 is small enough.

Lemma 19. Let $z_0 \in \partial M$, $t_0 \in (0, \tau_{\text{exit}}(\nu(z_0)))$ and $F(z_0, t_0) = \{U', V(\cdot), t(\cdot)\}$ be a family of focusing directions around (z_0, t_0) . If

$$t_0 < \tau_{\text{cut}}(z_0, \nu(z_0)) =: \tau_M(z_0),$$

then there exists a neighborhood $U \subset U'$ of z_0 such that (5) holds true.

For the proof of the lemma we need the following auxiliary result.

Lemma 20. Let $z_0 \in \partial M$, $t_0 > 0$, and the family of focusing directions $F(z_0, t_0) = \{U, V(\cdot), t(\cdot)\}$ around (z_0, t_0) be as in Lemma 19. Let γ be some geodesic that intersects $\gamma_{V(z_0)}$ at r_0 transversely. If there exist functions $r, \rho : U \rightarrow \mathbb{R}$ for which hold

$$\begin{aligned} \rho(z_0) &= 0, \quad r(z_0) = r_0, \quad \gamma(\rho(z)) = \gamma_{V(z)}(r(z)), \\ 0 \leq r(z) &\leq r_1 < \tau_M(z_0), \quad |\rho(z)| \leq \rho_1 < \text{inj}(M). \end{aligned}$$

Then $t_0 = r_0$.

Proof. The proof consists of several steps.

(I) Since $F(z_0, t_0)$ is a family of focusing directions, equation (3) implies the existence of functions $s, \hat{s} : U \rightarrow [0, \infty)$ that satisfy

$$\gamma_{V(z)}(s(z)) = \gamma_{V(z_0)}(\hat{s}(z)), \quad s(z) + \hat{s}(z) = t(z) + t_0.$$

Since the vector field V is continuous, equation (4) and Lemma 16 imply

$$s(z) \rightarrow t_0, \quad \hat{s}(z) \rightarrow t_0 \quad \text{as } z \rightarrow z_0.$$

Therefore $s(z_0) = \hat{s}(z_0) = t_0$, and by an analogous proof to one given in [Kurylev et al. 2010, Lemma 2.8] we show that the functions s, \hat{s} are smooth near z_0 and satisfy

$$ds(z_0) = d\hat{s}(z_0) = 0.$$

(II) Let $W \subset \partial M$ be a neighborhood of z_0 where the functions s, \hat{s} are smooth. We consider a map

$$E : W \rightarrow SM, \quad E(z) := -\phi_{s(z)}(V(z)).$$

We begin by studying the differential $dE|_{z_0} : T_{z_0} \partial M \rightarrow T_{(x,\eta)} SM, (x, \eta) := E(z_0)$. Recall that the tangent bundle $T(SM)$ has a canonical decomposition to horizontal $H(SM)$ and vertical $V(SM)$ subbundles. We denote the projections from $T(SM)$ to these bundles by P_H and P_V respectively. We note first that

$$x(z) := (\pi \circ E)(z) = \gamma_{V(z)}(s(z)) = \gamma_v(\hat{s}(z)).$$

Also a simple computation shows that after identifying $H_{(x,\eta)}(SM)$ to $T_x M$ we have $dx|_{z_0} = (P_H \circ (dE))|_{z_0}$. Therefore

$$(P_H \circ (dE))|_{z_0} = d[\gamma_{V(z)}(s(z))] \Big|_{z=z_0} = d[\gamma_v(\hat{s}(z))] \Big|_{z=z_0} = \dot{\gamma}_v(t_0) \otimes d\hat{s}(z_0) = 0$$

implies that

$$dE|_{z_0} : T_{z_0} \partial M \rightarrow T_{(x,\eta)} SM = H_{(x,\eta)}(SM) \oplus V_{(x,\eta)}(SM), \quad dE|_{z_0} v = (0, \Theta v),$$

where $\Theta : T_{z_0} \partial M \rightarrow V_{(x,\eta)}(SM)$ is a linear map. Recall that $V_{(x,\eta)}(SM)$ is isomorphic to $T_\eta(S_x M)$, which is $(n-1)$ -dimensional, as is $T_{z_0} \partial M$. Therefore to show that Θ is a linear isomorphism it suffices to prove that it is injective. We define a map

$$G(z) := \exp_{x(z)}(s(z)E(z)) = z \quad \text{for all } z \in W.$$

Thus the differential dG at x_0 is an identity operator on $T_{z_0} \partial M$. Since $s(z_0) = t_0$ is less than the cut distance $\tau_{\text{cut}}(z_0, v(z_0))$ and $ds(z)|_{z=z_0} = dx(z)|_{z=z_0} = 0$, we have

$$dG|_{z=z_0} v = d(\exp_x)|_{t_0 E(z_0)} t_0 dE|_{z=z_0} v = d(\exp_x)|_{t_0 E(z_0)} t_0 \Theta v = v.$$

This implies that Θ is an injection.

(III) Now we prove that the function r given in the claim of this lemma is continuous at z_0 . If this is not true, there exist $\epsilon > 0$ and a sequence $z_k \in \partial M$ which converges to z_0 but for which hold

$$|r_k - r_0| > \epsilon, \quad r_k := r(z_k).$$

Since functions r and ρ are bounded we can without loss of generality assume that $r_k \rightarrow r' < \tau_M(z_0)$, $|r' - r_0| \geq \epsilon$, $\rho_k := \rho(z_k) \rightarrow \rho' \in \mathbb{R}$ such that $|\rho'| < \text{inj}(M)$. Thus

$$\gamma(\rho') = \lim_{k \rightarrow \infty} \gamma(\rho_k) = \lim_{k \rightarrow \infty} \gamma_{V(z_k)}(r_k) = \gamma_v(r').$$

Since $r_0, r' < \tau_M(z_0)$ we have that $x := \gamma_v(r_0) = \gamma(0)$ and $x' := \gamma_v(r') = \gamma(\rho')$ are two different points where γ and γ_v intersect. Since $r_0, r' < \tau_M(z_0)$, and $|\rho'| < \text{inj}(M)$, there are two different distance-minimizing geodesics connecting x to x' , which is not possible.

(IV) Let us then assume that $r_0 < t_0$. We study a map

$$\Phi : U \times \mathbb{R} \rightarrow M, \quad \Phi(z, \lambda) = \exp_z(\lambda V(z)),$$

and show that it is a local diffeomorphism near (z_0, r_0) . Since $0 < t_0 - r_0 < \tau_M(z_0)$ and the Finsler function F is reversible, the map \exp_{x_0} is a local diffeomorphism near $(t_0 - r_0)\eta$, where $x_0 = \gamma_v(t_0 - r_0)$, $\eta_0 = -\dot{\gamma}_v(t_0 - r_0)$. Thus

$$d\exp_{x_0}|_{(t_0-r_0)\eta_0} : T_{(t_0-r_0)\eta}(T_{x_0}M) \rightarrow T_{z_0}M$$

is a linear isomorphism. Next we note that

$$\Phi(z, \lambda) = \gamma_{E(z)}(s(z) - \lambda) = \exp_{x(z)}((s(z) - \lambda)E(z)),$$

where E and $s(\cdot)$ are the same maps as in parts (I) and (II) of the proof. Since the differentials of x and s vanish at z_0 we have

$$\begin{aligned} d_\lambda \Phi|_{(z_0, r_0)} &= -d\exp_{x_0}|_{(t_0-r_0)E(z_0)} E(z_0) := -AE(z_0), \\ d_z \Phi|_{(z_0, r_0)} &= (t_0 - r_0)AdE|_{z_0}. \end{aligned}$$

Therefore for any $(v, \lambda) \in T_{z_0} \partial M \times \mathbb{R}$ it holds that

$$d\Phi|_{(z_0, r_0)}(v, \lambda) = A((t_0 - r_0)\Theta v - \lambda E(z_0)).$$

Since A is invertible it holds that

$$d\Phi|_{(z_0, r_0)}(v, \lambda) = 0 \quad \text{if and only if} \quad (t_0 - r_0)\Theta v - \lambda E(z_0) = 0.$$

However, this can only happen if $(v, \lambda) = 0$ as Θ is injective and

$$\Theta v \in T_{E(z_0)}S_{x_0}M \quad \text{implies} \quad g_{E(z_0)}(E(z_0), \Theta v) = 0.$$

Due to the inverse function theorem, Φ is a local diffeomorphism near (z_0, r_0) .

(V) Let Σ be an $(n-1)$ -dimensional surface in M such that γ is a curve on Σ and γ_v is transverse to Σ at r_0 . Due to the implicit function theorem, there exists a smooth function $\hat{r} : U \rightarrow \mathbb{R}$ so that

$$\Phi(z, \lambda) \in \Sigma \quad \text{if and only if} \quad \lambda = \hat{r}(z) \text{ for all } z \in U \quad \text{and} \quad \hat{r}(z_0) = r_0.$$

Since $r(\cdot)$ is continuous and satisfies $\Phi(z, r(z)) = \gamma(\rho(z)) \in \Sigma$, it holds that near z_0 the functions r and \hat{r} coincide. Therefore the map

$$\tilde{\Phi} : U \rightarrow \Sigma, \quad z \mapsto \Phi(z, r(z)),$$

is smooth and its image is contained in the image of γ . However, we have proven that $\tilde{\Phi}$ is a local diffeomorphism near z_0 . Thus we arrive at a contradiction since $\tilde{\Phi}$ maps an $(n-1)$ -dimensional surface $(n-1 \geq 2)$ onto a 1-dimensional surface. Thus $r_0 < t_0$ is false. By an analogous argument we can prove that $r_0 > t_0$ is also false and therefore it must hold that $r_0 = t_0$. □

We are ready to present the proof for [Lemma 19](#).

Proof of Lemma 19. The proof is an adaptation of [[Kurylev et al. 2010](#), Proof of Theorem 2.6, Step 4] where one uses [Lemmas 16](#) and [20](#). □

5.3. Boundary distance functions. We start with giving a stronger formulation of [Lemma 20](#).

Lemma 21. *Let $z_0 \in \partial M$, $0 < t_0 < \tau_M(z_0)$, and $F(z_0, t_0) = \{U, V(\cdot), t(\cdot)\}$ be a family of focusing directions around (z_0, t_0) . Let γ be some geodesic that intersects every geodesic of $F(z_0, t_0)$ in the sense that*

$$\gamma(\rho(z)) = \gamma_{V(z)}(r(z))$$

for some functions $\rho, r : U \rightarrow \mathbb{R}$ satisfying

$$0 \leq r(z) \leq r_1 < \tau_M(z_0), \quad |\rho(z)| < L \quad \text{for some } L > 0.$$

If in addition $h(z) := r(z) + \rho(z)$ is continuous then for any $z \in \partial M$ which is close to z_0 it holds that

$$\gamma(h(z_0) - t_0) = \gamma_{V(z_0)}(t(z)).$$

That is, all geodesics of $F(z_0, t_0)$ meet at the same point $\gamma(h(z_0) - t_0)$.

Proof. If γ and $\gamma_{V(z_0)}$ are the same geodesic, the result follows from [Lemma 19](#).

If we assume that γ is a different geodesic than $\gamma_{V(z_0)}$, it then holds due to the definition of the injectivity radius of M that the geodesic segments $\gamma([-L, L])$ and $\gamma_{V(z_0)}([0, r_1])$ can intersect at most finitely many times. Thus there exists $N \in \mathbb{N}$ such that $\gamma(\rho) = \gamma_{V(z_0)}(r)$ for $(\rho, r) \in [-L, L] \times (0, r_1)$ if and only if $(\rho, r) = (\rho_0^k, r_0^k)$ for some $\rho_0^k \in \{\rho_0^1, \dots, \rho_0^N\} \subset [-L, L]$, and $r_0^k \in \{r_0^1, \dots, r_0^N\} \subset (0, r_1)$.

Let $0 < \epsilon < \frac{1}{2} \text{inj}(M)$. We claim that there exists $R_0 > 0$ such that for all $0 < R < R_0$ and

$$z \in U(R) := B(z_0, R) \cap \partial M \quad \text{it holds that} \quad \min_{j \in \{1, \dots, N\}} |\rho(z) - \rho_0^j| < \epsilon. \quad (6)$$

If this is not true, the boundedness of the functions $r(\cdot)$ and $\rho(\cdot)$ imply the existence of a sequence $(z_k)_{k=1}^\infty \subset \partial M$ that converges to z_0 and which satisfies

$$\lim_{k \rightarrow \infty} r(z_k) = \tilde{r} \in [0, r_1], \quad \lim_{k \rightarrow \infty} \rho(z_k) = \hat{\rho} \in [-L, L] \setminus \{\rho_0^1, \dots, \rho_0^N\}.$$

Therefore we arrive at the contradiction $\gamma(\hat{\rho}) = \gamma_{V(z_0)}(\hat{r})$.

We set

$$W_j(R) := \{z \in U(R) : \text{there exists } r \in [0, r_1], \text{ there exists } \rho \in [-L, L] \\ \text{such that } \gamma_{V(z)}(r) = \gamma(\rho), r + \rho = h(z), |\rho - \rho_0^j| \leq \epsilon, \quad j \in \{1, \dots, N\},$$

and show that these sets are relatively closed in $U(R)$. Choose a sequence $(z_k)_{k=1}^\infty \subset W_j(R)$ that converges to $z \in U(R)$. For any $k \in \mathbb{N}$ we choose $r_k \in [0, r_1]$ and $\rho_k \in [\rho_0^j - \epsilon, \rho_0^j + \epsilon]$ for which

$$\gamma_{V(z_k)}(r_k) = \gamma(\rho_k) \quad \text{and} \quad r_k + \rho_k = h(z_k).$$

After choosing a subsequence of $(z_k)_{k=1}^\infty$, we may without loss of generality assume that $\lim_{k \rightarrow \infty} r_k = \tilde{r} \in [0, r_1]$, $\lim_{k \rightarrow \infty} \rho_k = \tilde{\rho} \in [\rho_0^j - \epsilon, \rho_0^j + \epsilon]$,

$$\gamma_{V(z)}(\tilde{r}) = \gamma(\tilde{\rho}) \quad \text{and} \quad h(z) = \tilde{r} + \tilde{\rho}.$$

Here we used the continuity of the geodesic flow and the function $h(\cdot)$ to obtain the last two equations. Thus we have verified that $z \in W_j(R)$. Therefore $W_j(R)$ is closed and measurable.

Equation (6) implies that

$$U(R) = \bigcup_{j=1}^N W_j(R).$$

Thus there exists $j \in \{1, \dots, N\}$ for which $W_j(R)$ has a strictly positive $(n-1)$ -dimensional measure. Choose $k \in \{1, \dots, N\}$ and suppose that $r_0^k \neq t_0$. By following the proof of Lemma 20 we show that after choosing a smaller $R > 0$ there exist $0 < T < \frac{1}{2} \text{inj}(M)$, an $(n-1)$ -dimensional surface $\Sigma \subset M$ and a diffeomorphism

$$\tilde{\Phi} : U(R) \rightarrow \Sigma \quad \text{for which } \tilde{\Phi}(z) \subset \gamma([\rho_0^k - T, \rho_0^k + T]), \text{ if } z \in W_k(R).$$

Thus the $(n-1)$ -dimensional measure of $W_k(R)$ is zero. We have proven that the set $K \subset \{1, \dots, N\}$ that contains all those $k \in \{1, \dots, N\}$ for which $r_0^k = t_0$, is not empty and moreover

$$U(R) \setminus \left(\bigcup_{k \in K} W_k(R) \right) \text{ is of measure zero.}$$

Therefore $\bigcup_{k \in K} W_k(R)$ is dense in $U(R)$. Since $\epsilon > 0$ was arbitrary, we can choose $k \in K$, sequences $(z_j)_{j=1}^\infty \subset W_k(R)$, $(r_j)_{j=1}^\infty$, $(\rho_j)_{j=1}^\infty \subset \mathbb{R}$, such that $z_j \rightarrow z_0$, $\rho_j \rightarrow \rho_0^k$, $r_j \rightarrow \tilde{r} \in [0, r_1]$ and $\gamma(\rho_j) = \gamma_{V(z_j)}(r_j)$ for every $j \in \mathbb{N}$. Since $r_0^k = t_0$, the continuity of geodesic flow implies

$$\gamma_{V(z_0)}(r_0^k) = \gamma(\rho_0^k) = \gamma_{V(z_0)}(\tilde{r}).$$

Thus we must have $t_0 = \tilde{r}$. Since the function $h(\cdot)$ is continuous, we also have $\rho_0^k = h(z_0) - t_0$ and moreover $\gamma(h(z_0) - t_0) = \gamma_{V(z_0)}(t_0)$. This and Lemma 19 complete the proof. \square

Lemma 22. *The broken scattering relation determines the boundary cut distance function $\tau_{\partial M}$.*

Proof. Let $z_0 \in \partial M$. By the definition of $\tau_{\partial M}(z_0)$, it holds that for $t_0 \in (0, \tau_{\partial M}(z_0))$ the point z_0 is the closest boundary point to $x_0 := \gamma_{v(z_0)}(t_0)$. However, if $t_0 > \tau_{\partial M}(z_0)$ then there exists $w \in \partial M$ that is closer to x_0 than z_0 in the sense that

$$\text{there exists } s < t_0 \text{ such that } \gamma_{v(w)}(s) = x_0.$$

By Lemma 17 we know that $\tau_{\text{cut}}(z_0, v(z_0)) > \tau_{\partial M}(z_0)$. If $t_0 \in (\tau_{\partial M}(z_0), \tau_M(z_0))$ then according to the discussion at the beginning of Section 5.1, there exist $w \in \partial M$ and a family of focusing directions $F(z_0, t_0) := \{U, V(\cdot), t(\cdot)\}$ around (z_0, t_0) such that

$$v(w)R_{s+t(z)}V(z) \quad \text{and} \quad s < t(z) \quad \text{for all } z \in U. \tag{7}$$

First we note that the broken scattering relation determines the exit time function since for any $v \in \overline{\partial_{\text{in}} SM}$

$$\tau_{\text{exit}}(v) = \sup\{t \geq 0 : vR_{2t}v\}.$$

We claim that

$$\tau_{\partial M}(z_0) = \inf\{t_0 \in (0, \tau_{\text{exit}}(v(z_0))) : \text{there exist } F(z_0, t_0), s < t_0, \text{ and } w \in \partial M \setminus \{z_0\} \text{ such that (7) is valid}\}.$$

For the proof of this equation, see in [Kurylev et al. 2010, Lemma 2.10]. \square

We recall that the exponential map at the boundary is

$$\exp_{\partial M} : \partial M \times [0, \infty) \rightarrow M, \quad \exp_{\partial M}(z, t) = \gamma_{\nu(z)}(t).$$

Proposition 23. *Let $x \in M$. The broken scattering relation determines the set of boundary distance functions*

$$r_x : \partial M \rightarrow \mathbb{R}, \quad r_x(z) = d(x, z), \quad z \in \partial M, x \in M.$$

Proof. By the definition of boundary cut distance function we have that

$$M = \{\exp_{\partial M}(z, t) \in M : z \in \partial M, t \in [0, \tau_{\partial M}(z)]\}.$$

Therefore $\exp_{\partial M}^{-1} M \subset (\partial M \times \mathbb{R})$ is a way to represent M , although points in the boundary cut locus

$$\omega_{\partial M} := \{x \in M : x = \exp_{\partial M}(z, \tau_{\partial M}(z)), z \in \partial M\}$$

may have several representatives. As we are interested in determining the boundary distance functions, the possible ambiguity does not concern us since it is proven in [Hoop et al. 2019, Proposition 3.1] that

$$r_x = r_y \quad \text{if and only if} \quad x = y \in M.$$

Let us fix $z_0 \in \partial M$ and $t_0 \in [0, \tau_{\partial M}(z_0)]$. Then we choose $w \in \partial M$. As ∂M is strictly convex, any distance-minimizing curve from w to $x := \exp_{\partial M}(z_0, t_0)$ is a geodesic $\gamma_{w, \eta} : [0, s] \rightarrow M$, where $s = d(w, x)$ and $\eta \in S_w M$. Due to Lemma 17 there exists a family of focusing directions $F(z_0, t_0) = \{U, V(\cdot), t(\cdot)\}$ such that $V(z)R_{s+t(z)}\eta$ for any $z \in U$. We set

$$S := \{s > 0 : \text{there exist } \eta \in S_w M \text{ and } F(z_0, t_0) \text{ such that } V(z)R_{s+t(z)}\eta \text{ holds}\}$$

and claim that

$$d(x, w) = \inf S.$$

The proof of this claim is an adaptation of the proof of an analogous statement in [Kurylev et al. 2010, Theorem 2.13]. \square

In the following lemma we give an invariant definition for families of focusing directions using the invariant definition of broken scattering relation given in Theorem 4. The claim of the lemma is a direct implication of the Definition 18.

Lemma 24. *Let (M_i, F_i) be two compact Finsler manifolds that satisfy the conditions of Theorem 4. Let $z_0 \in \partial M_1$ and $t_0 \in (0, \tau_{\text{exit}}(\nu(z_0)))$. Then $\mathcal{F}(z_0, t_0) := \{U, V(\cdot), t(\cdot)\}$ is a family of focusing directions around (z_0, t_0) if and only if $\tilde{\mathcal{F}}(z_0, t_0) := \{\tilde{U}, \tilde{V}(\cdot), \tilde{t}(\cdot)\}$ is a family of focusing directions around $(\xi(z_0), t_0)$, where*

$$\tilde{U} := \xi(U) \subset \partial M_2, \quad \tilde{V} := \Xi \circ V \circ \xi^{-1}, \quad \text{and} \quad \tilde{t} := t \circ \xi^{-1}.$$

We are ready to prove Proposition 7.

Proof of Proposition 7. Let $x \in \text{int } M_1$. Choose $z_0 \in \partial M_1$ and $t_0 \in (0, \tau_{\partial M_1}(z_0)]$ such that $x = \exp_{\partial M_1}(z_0, t_0)$. Due to Lemmas 22 and 24, it holds that $\tau_{\partial M_2}(\xi(z_0)) = \tau_{\partial M_1}(z_0)$ and therefore $\tilde{x} := \exp_{\partial M_2}(\xi(z_0), t_0) \in \text{int } M_2$. We aim to verify

$$d_1(z, x) = d_2(\xi(z), \tilde{x}) \quad \text{for all } z \in \partial M_1. \quad (8)$$

This implies the left-hand side inclusion in (1). By reversing the roles of M_1 and M_2 , the analogous argument verifies the right-hand side inclusion in (1). Choose $z \in \partial M_1$. We define $s_1 := d_1(x, z)$ and $s_2 := d_2(\tilde{x}, \xi(z))$. Since ∂M_1 is strictly convex there exists a distance-minimizing geodesic $\gamma_{z,\eta}$ of M_1 from z to x . Choose a family of focusing directions $\mathcal{F}(z_0, t_0) = \{U, V(\cdot), t(\cdot)\}$ such that $V(z)R_{s_1+t(z)}^{(1)}\eta$ for every $z \in U$. Then Lemma 24 implies that $\tilde{\mathcal{F}}(z_0, t_0) = \{\tilde{U}, \tilde{V}(\cdot), \tilde{t}(\cdot)\}$ is a focusing family at $(\xi(z_0), t_0)$ that satisfies $\tilde{V}(w)R_{s_1+\tilde{t}(w)}^{(2)}\Xi\eta$ for every $w \in \tilde{U}$. The proof of Proposition 23 implies $s_2 \leq s_1$. After reversing the roles of x and \tilde{x} , we use the analogous argument to verify $s_1 \leq s_2$. We have proven $s_1 = s_2$. Since $z \in M_1$ was arbitrary, the equation (8) is valid. \square

Appendix: Fourier integral operators and annihilators

We consider $v, w \in \partial SM$ and introduce their duals, $\tilde{v}, \tilde{w} \in \partial(S^*M)$, via the Legendre transform [Shen 2001, Formula (3.4)]. Their tangential components in $T^*\partial M$ are written as $\tilde{v}_\partial, \tilde{w}_\partial$. Instead of the geodesic flow, ϕ_t , we consider the cogeodesic or Hamiltonian flow, $\tilde{\phi}_t$ on the cosphere bundle and the canonical projection $\tilde{\pi} : S^*M \rightarrow M$. We note that the broken scattering relation, given in Definition 3, is closely related to the canonical relation

$$\Lambda := \{(\tilde{\pi}(\tilde{v}), \tilde{\pi}(\tilde{w}), t, \tilde{v}_\partial, \tilde{w}_\partial, \tau); (x, \xi) \mid \tilde{\pi}(\tilde{\phi}_{t_1}(\tilde{v})) = \tilde{\pi}(\tilde{\phi}_{t_2}(\tilde{w})) = x, t = t_1 + t_2, \tau = F^*(\tilde{\phi}_{t_1}(\tilde{v})) = F^*(\tilde{\phi}_{t_2}(\tilde{w})) = 1, \xi = \tilde{\phi}_{t_1}(\tilde{v}) + \tilde{\phi}_{t_2}(\tilde{w})\}.$$

We write $Y = \partial M \times \partial M \times (0, T)$ and $X = M$. Then $\Lambda \subset T^*Y \times T^*X$. If the Finsler metric on ∂TM is known, as we assume in Theorem 4, the vectors $\tilde{v}_\partial, \tilde{w}_\partial$ determine \tilde{v}, \tilde{w} , respectively. The canonical relation connects an element (x, ξ) in the wavefront set of scatterers in T^*X to an element in the wavefront set of scattered waves generated by sources, and detected by receivers, in ∂M in T^*Y . That is, there is a Fourier integral operator (FIO) that maps scatterers to scattered waves restricted to ∂M and propagates singularities according to the mentioned canonical relation Λ .

The Bolker condition states that the natural projection from Λ to T^*Y is injective. If the Bolker condition is satisfied, the range of the FIO can be characterized by pseudodifferential annihilators [de Hoop and Uhlmann 2006; Guillemin 1985], which yields the canonical relation. Thus, within the framework of the results of this paper, vanishing annihilators determine the underlying reversible Finsler manifold if both the Bolker condition and the foliation condition of Definition 1 hold.

Although the Bolker condition should be fairly easy to satisfy, it does not follow from the foliation condition. To see this, we consider the Euclidean unit disc M with a radial metric $c(r)e, r = |x|$. This manifold can satisfy the Herglotz condition,

$$\frac{d}{dr} \left(\frac{r}{c(r)} \right) > 0$$

and have a geodesic whose opening angle is between π and 2π . Informally, this corresponds to a geodesic going around the center. Such a geodesic will meet its reflection across the center two times at some points $p, q \in \text{int } M$. If we remove a segment of the geodesic between the second intersection point q and the nearest endpoint at the boundary and the mirror image of this segment, we obtain two broken rays c_1




Figure 1. Here c_1 is the broken ray starting with initial velocity v , splitting at p and exiting with velocity $-w$; c_2 is the broken ray with same boundary conditions $\{v, -w\}$ that splits at q .

and c_2 with exactly the same total length $t \in \mathbb{R}$ and boundary data $v, w \in \partial_{\text{in}} SM$. This situation violates the Bolker condition. We illustrate this setup in [Figure 1](#).

Acknowledgements

De Hoop was supported by the Simons Foundation under the MATH+X program, the National Science Foundation under grant DMS-1815143, and the corporate members of the Geo-Mathematical Imaging Group at Rice University. Ilmavirta was supported by the Academy of Finland (project 295853). Lassas was supported by Academy of Finland (projects 284715 and 303754). Saksala was supported by the Simons Foundation under the MATH+X program and the corporate members of the Geo-Mathematical Imaging Group at Rice University.

Part of this work was carried out during the visit of Ilmavirta and Saksala to the University of Washington and they are grateful for Prof. Gunther Uhlmann for hospitality and support.

We wish to express our gratitude for the anonymous referees. Their invaluable comments improved the manuscript considerably.

References

- [Bal et al. 2015] G. Bal, F. Monard, and G. Uhlmann, “Reconstruction of a fully anisotropic elasticity tensor from knowledge of displacement fields”, *SIAM J. Appl. Math.* **75**:5 (2015), 2214–2231. [MR](#) [Zbl](#)
- [Bao et al. 2018] G. Bao, G. Hu, J. Sun, and T. Yin, “Direct and inverse elastic scattering from anisotropic media”, *J. Math. Pures Appl.* (9) **117** (2018), 263–301. [MR](#) [Zbl](#)
- [Barceló et al. 2018] J. A. Barceló, M. Folch-Gabayet, S. Pérez-Esteva, A. Ruiz, and M. C. Vilela, “Uniqueness for inverse elastic medium problems”, *SIAM J. Math. Anal.* **50**:4 (2018), 3939–3962. [MR](#) [Zbl](#)
- [Bartolo et al. 2011] R. Bartolo, E. Caponio, A. V. Germinario, and M. Sánchez, “Convex domains of Finsler and Riemannian manifolds”, *Calc. Var. Partial Differential Equations* **40**:3-4 (2011), 335–356. [MR](#) [Zbl](#)
- [Beretta et al. 2014] E. Beretta, E. Francini, and S. Vessella, “Uniqueness and Lipschitz stability for the identification of Lamé parameters from boundary measurements”, *Inverse Probl. Imaging* **8**:3 (2014), 611–644. [MR](#) [Zbl](#)
- [Burago and Ivanov 2010] D. Burago and S. Ivanov, “Boundary rigidity and filling volume minimality of metrics close to a flat one”, *Ann. of Math.* (2) **171**:2 (2010), 1183–1211. [MR](#) [Zbl](#)
- [do Carmo 1992] M. P. a. do Carmo, *Riemannian geometry*, Birkhäuser, Boston, MA, 1992. [MR](#) [Zbl](#)

- [Croke 1991] C. B. Croke, “Rigidity and the distance between boundary points”, *J. Differential Geom.* **33**:2 (1991), 445–464. [MR](#) [Zbl](#)
- [Croke and Kleiner 1994] C. B. Croke and B. Kleiner, “Conjugacy and rigidity for manifolds with a parallel vector field”, *J. Differential Geom.* **39**:3 (1994), 659–680. [MR](#) [Zbl](#)
- [Duistermaat 1996] J. J. Duistermaat, *Fourier integral operators*, Progress in Mathematics **130**, Birkhäuser, Boston, MA, 1996. [MR](#) [Zbl](#)
- [Dziewonski and Anderson 1981] A. M. Dziewonski and D. L. Anderson, “Preliminary reference Earth model”, *Phys. Earth Planet. Int.* **25**:4 (1981), 297–356.
- [Greenleaf and Uhlmann 1993] A. Greenleaf and G. Uhlmann, “Recovering singularities of a potential from singularities of scattering data”, *Comm. Math. Phys.* **157**:3 (1993), 549–572. [MR](#) [Zbl](#)
- [Greenleaf et al. 2003] A. Greenleaf, M. Lassas, and G. Uhlmann, “On nonuniqueness for Calderón’s inverse problem”, *Math. Res. Lett.* **10**:5-6 (2003), 685–693. [MR](#) [Zbl](#)
- [Greenleaf et al. 2009] A. Greenleaf, Y. Kurylev, M. Lassas, and G. Uhlmann, “Invisibility and inverse problems”, *Bull. Amer. Math. Soc. (N.S.)* **46**:1 (2009), 55–97. [MR](#) [Zbl](#)
- [Gromov 1983] M. Gromov, “Filling Riemannian manifolds”, *J. Differential Geom.* **18**:1 (1983), 1–147. [MR](#) [Zbl](#)
- [Guillemin 1985] V. Guillemin, “On some results of Gelfand in integral geometry”, pp. 149–155 in *Pseudodifferential operators and applications* (Notre Dame, IN, 1984), edited by F. Trèves, Proc. Sympos. Pure Math. **43**, Amer. Math. Soc., Providence, RI, 1985. [MR](#) [Zbl](#)
- [Herglotz 1905] G. Herglotz, “Über die Elastizität der Erde bei Berücksichtigung ihrer variablen Dichte”, *Z. Math. Phys.* **52** (1905), 275–299. [JFM](#)
- [de Hoop and Saksala 2019] M. V. de Hoop and T. Saksala, “Inverse problem of travel time difference functions on a compact Riemannian manifold with boundary”, *J. Geom. Anal.* **29**:4 (2019), 3308–3327. [MR](#) [Zbl](#)
- [de Hoop and Uhlmann 2006] M. V. de Hoop and G. Uhlmann, “Characterization and ‘source-receiver’ continuation of seismic reflection data”, *Comm. Math. Phys.* **263**:1 (2006), 1–19. [MR](#) [Zbl](#)
- [Hoop et al. 2019] M. V. D. Hoop, J. Ilmavirta, M. Lassas, and T. Saksala, “Determination of a compact Finsler manifold from its boundary distance map and an inverse problem in elasticity”, preprint, 2019. To appear in *Comm. Anal. Geom.* [arXiv](#)
- [Hu et al. 2014] G. Hu, J. Li, and H. Liu, “Recovering complex elastic scatterers by a single far-field pattern”, *J. Differential Equations* **257**:2 (2014), 469–489. [MR](#) [Zbl](#)
- [Ivanov 2013] S. Ivanov, “Local monotonicity of Riemannian and Finsler volume with respect to boundary distances”, *Geom. Dedicata* **164** (2013), 83–96. [MR](#) [Zbl](#)
- [Ivanov 2020] S. Ivanov, “Distance difference representations of Riemannian manifolds”, *Geom. Dedicata* **207** (2020), 167–192. [MR](#) [Zbl](#)
- [Katchalov et al. 2001] A. Katchalov, Y. Kurylev, and M. Lassas, *Inverse boundary spectral problems*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics **123**, Chapman & Hall/CRC, Boca Raton, FL, 2001. [MR](#) [Zbl](#)
- [Kurylev 1997] Y. Kurylev, “Multidimensional Gelfand inverse problem and boundary distance map”, pp. 1–15 in *Inverse problems related with geometry*, edited by H. Soga, 1997. [Zbl](#)
- [Kurylev et al. 2010] Y. Kurylev, M. Lassas, and G. Uhlmann, “Rigidity of broken geodesic flow and inverse problems”, *Amer. J. Math.* **132**:2 (2010), 529–562. [MR](#) [Zbl](#)
- [Lassas and Saksala 2019] M. Lassas and T. Saksala, “Determination of a Riemannian manifold from the distance difference functions”, *Asian J. Math.* **23**:2 (2019), 173–199. [MR](#) [Zbl](#)
- [Lassas et al. 2003] M. Lassas, V. Sharafutdinov, and G. Uhlmann, “Semiglobal boundary rigidity for Riemannian metrics”, *Math. Ann.* **325**:4 (2003), 767–793. [MR](#) [Zbl](#)
- [Lassas et al. 2018] M. Lassas, T. Saksala, and H. Zhou, “Reconstruction of a compact manifold from the scattering data of internal sources”, *Inverse Probl. Imaging* **12**:4 (2018), 993–1031. [MR](#) [Zbl](#)
- [Liu and Xiao 2017] H. Liu and J. Xiao, “Decoupling elastic waves and its applications”, *J. Differential Equations* **263**:8 (2017), 4442–4480. [MR](#) [Zbl](#)

- [Mazzucato and Rachele 2007] A. L. Mazzucato and L. V. Rachele, “On uniqueness in the inverse problem for transversely isotropic elastic media with a disjoint wave mode”, *Wave Motion* **44**:7-8 (2007), 605–625. [MR](#) [Zbl](#)
- [Michel 1981] R. Michel, “Sur la rigidité imposée par la longueur des géodésiques”, *Invent. Math.* **65**:1 (1981), 71–83. [MR](#) [Zbl](#)
- [Michel 1994] R. Michel, “Restriction de la distance géodésique à un arc et rigidité”, *Bull. Soc. Math. France* **122**:3 (1994), 435–442. [MR](#) [Zbl](#)
- [Muhometov 1981] R. G. Muhometov, “On a problem of reconstructing Riemannian metrics”, *Sibirsk. Mat. Zh.* **22**:3 (1981), 119–135. In Russian; translated in *Sib. Math. J.* **22**:3 (1981), 420–433. [MR](#)
- [Nakamura and Uhlmann 2003] G. Nakamura and G. Uhlmann, “Erratum to ‘Global uniqueness for an inverse boundary value problem arising in elasticity’”, *Invent. Math.* **152**:1 (2003), 205–207. [MR](#)
- [Oksanen 2011] L. Oksanen, “Solving an inverse problem for the wave equation by using a minimization algorithm and time-reversed measurements”, *Inverse Probl. Imaging* **5**:3 (2011), 731–744. [MR](#) [Zbl](#)
- [Paternain et al. 2019] G. P. Paternain, M. Salo, G. Uhlmann, and H. Zhou, “The geodesic X-ray transform with matrix weights”, *Amer. J. Math.* **141**:6 (2019), 1707–1750. [MR](#) [Zbl](#)
- [Pestov and Uhlmann 2005] L. Pestov and G. Uhlmann, “Two dimensional compact simple Riemannian manifolds are boundary distance rigid”, *Ann. of Math. (2)* **161**:2 (2005), 1093–1110. [MR](#) [Zbl](#)
- [Shen 2001] Z. Shen, *Lectures on Finsler geometry*, World Scientific, Singapore, 2001. [MR](#) [Zbl](#)
- [Sklan et al. 2018] S. R. Sklan, R. Y. S. Pak, and B. Li, “Seismic invisibility: elastic wave cloaking via symmetrized transformation media”, *New J. of Phys.* **20**:6 (2018), art. id. 063013.
- [Stefanov and Uhlmann 2005] P. Stefanov and G. Uhlmann, “Boundary rigidity and stability for generic simple metrics”, *J. Amer. Math. Soc.* **18**:4 (2005), 975–1003. [MR](#) [Zbl](#)
- [Stefanov and Uhlmann 2018] P. Stefanov and G. Uhlmann, “Microlocal analysis and integral geometry”, book preprint, 2018, available at <https://www.math.purdue.edu/~stefanop/publications/book.pdf>.
- [Stefanov et al. 2016] P. Stefanov, G. Uhlmann, and A. Vasy, “Boundary rigidity with partial data”, *J. Amer. Math. Soc.* **29**:2 (2016), 299–332. [MR](#) [Zbl](#)
- [Stefanov et al. 2021] P. Stefanov, G. Uhlmann, and A. Vasy, “Local and global boundary rigidity and the geodesic X-ray transform in the normal gauge”, *Ann. of Math. (2)* **194**:1 (2021), 1–95. [MR](#) [Zbl](#)
- [Uhlmann and Vasy 2016] G. Uhlmann and A. Vasy, “The inverse problem for the local geodesic ray transform”, *Invent. Math.* **205**:1 (2016), 83–120. [MR](#) [Zbl](#)

Received 21 May 2021. Revised 30 Aug 2021. Accepted 9 Oct 2021.

MAARTEN V. DE HOOP: mdehoop@rice.edu

Departments of Computational and Applied Mathematics, Earth Science, Rice University, Houston, TX, United States

JOONAS ILMAVIRTA: joonas.ilmavirta@jyu.fi

Department of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, Finland

MATTI LASSAS: matti.lassas@helsinki.fi

Department of Mathematics and Statistics, University of Helsinki, Helsinki, Finland

TEEMU SAKSALA: tssaksal@ncsu.edu

Department of Mathematics, North Carolina State University, Raleigh, NC, United States